

APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: METHODS OF INHIBITING FIBROSIS WITH A
SOMATOSTIN AGONIST

APPLICANT: MICHAEL D. CULLER AND PHILIP G. KASPRZYK

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL224702471US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

January 16, 2001

Date of Deposit

Signature

Samantha Bell
Typed or Printed Name of Person Signing Certificate

METHOD OF INHIBITING FIBROSIS WITH A SOMATOSTATIN AGONIST

BACKGROUND OF THE INVENTION

5
10
15
Tissue comprises organized cellular groups that are attached to an extracellular matrix and are surrounded by a network of blood vessels. Fibrosis is an abnormal accumulation of a collagen matrix following injury or inflammation which alters the structure and function of various tissues. Irrespective of location, the major pathology of fibrosis involves an excessive deposition of a collagen matrix which replaces the normal tissue at that site. Progressive fibrosis in the kidney, liver, lung, heart, bone or bone marrow, and skin is a major cause of death and suffering. See, e.g., Border, et al., New Engl. J. Med. 331:1286 (1994).

20
Development of fibrosis has been linked to the overexpression and over-production of TGF- β in numerous tissues and fibrotic disease states (see Border et al., N Engl J Med 1994, pp. 1286-92).

SUMMARY OF THE INVENTION

25
30
35
40
The present invention relates to a method of treating fibrosis in a patient (e.g., a mammal such as a human). The method includes the step of administering a therapeutically effective amount of somatostatin or a somatostatin agonist to said patient. The somatostatin or somatostatin agonist may be administered orally, topically, parenterally, e.g., administered intravenously, subcutaneously, or by implantation of a sustained release formulation. Fibrosis is the abnormal accumulation of an extracellular matrix (e.g., collagen) in tissue. The fibrosis, for example, may be located: in the kidney, for example, fibrosis as observed in glomerulonephritis, diabetic nephropathy), allograft rejection, and HIV nephropathy; in the liver, for example, cirrhosis, and veno-occlusive disease; in the lung, for example, idiopathic fibrosis; in the skin, for example, systemic sclerosis, keloids, scars, and eosinophilia-myalgia syndrome; in the central nervous system, for example,

-2-

intraocular fibrosis; in the cardiovascular system, for example, vascular restenosis; in the nose, for example, nasal polyposis; in bone or bone marrow; in an endocrine organ; and in the gastrointestinal system.

5 A fibrotic disorder may be induced by a number of causes including: chemotherapy, for example pulmonary fibrosis resulting from bleomycin, chlorambucil, cyclophosphamide, methotrexate, mustine, or procarbazine treatment; radiation
10 exposure whether accidental or purposeful as in radiation therapy, for example, interstitial lung disease (ILD) resulting from radiation; environmental or industrial factors or pollutants such as chemicals, fumes, metals, vapors, gases, etc. (e.g. ILD resulting from asbestos or coal dust); a drug or combination of drugs, for example, antibiotics (e.g.
15 penicillins, sulfonamides, etc.), cardiovascular drugs (e.g. hydralazine, beta blockers, etc.), CNS drugs (phenytoin, chlorpromazine, etc.) anti-inflammatory drugs (e.g. gold salts, phenylbutazone, etc.), etc. can cause ILD; an immune reaction disorder, for example, chronic graft-vs-host disease
20 with dermal fibrosis); disease states (e.g., aspiration pneumonia which is a known cause of ILD) which include parasite induced fibrosis; and wounds, for example, blunt trauma, surgical incisions, battlefield wounds, etc., as in penetrating injuries of the CNS.

25 In one aspect, this invention provides a method of inhibiting fibrosis in a patient, said method comprising administering a therapeutically effective amount of somatostatin or a somatostatin agonist to said patient; a method which is preferred of the foregoing method is wherein
30 said method comprises administering a therapeutically effective amount of a somatostatin agonist to said patient.

In another aspect, this invention provides a method of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective

amount of a somatostatin agonist wherein the fibrosis which is inhibited is in the:
kidney wherein the fibrotic disorder inhibited in the kidney is preferably glomerulonephritis, diabetic nephropathy, 5 allograft rejection or HIV nephropathy,
lung wherein the fibrotic disorder inhibited in the lung is preferably idiopathic fibrosis or autoimmune fibrosis,
liver wherein the fibrotic disorder inhibited in the liver is preferably cirrhosis or veno-occlusive disease,
10 skin wherein the fibrotic disorder inhibited in the skin is preferably systemic sclerosis, keloids, scars or eosinophilia-myalgia syndrome,
central nervous system wherein the fibrotic disorder inhibited in the central nervous system is preferably
15 intraocular fibrosis,
bone or bone marrow, cardiovascular system, an endocrine organ or gastrointestinal system. Each of the immediately foregoing methods is preferred.

In yet another aspect, this invention provides a method 20 of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective amount of a somatostatin agonist wherein the fibrosis is induced by chemotherapy and preferably the fibrosis inhibited is in the kidney, lung, liver, skin, central nervous system,
25 bone or bone marrow, cardiovascular system, an endocrine organ or gastrointestinal system. Each of the immediately foregoing methods is preferred.

In yet another aspect, this invention provides a method of inhibiting fibrosis in a patient which comprises 30 administering to the patient a therapeutically effective amount of a somatostatin agonist wherein the fibrosis is induced by radiation and preferably the fibrosis inhibited is in the kidney, lung, liver, skin, central nervous system, bone or bone marrow, cardiovascular system, an endocrine

organ or gastrointestinal system. Each of the immediately foregoing methods is preferred.

In yet another aspect, this invention provides a method of inhibiting fibrosis in a patient which comprises
5 administering to the patient a therapeutically effective amount of a somatostatin agonist wherein the fibrosis is induced by a drug or combination of drugs and preferably the fibrosis inhibited is in the kidney, lung, liver, skin, central nervous system, bone or bone marrow, cardiovascular
10 system, an endocrine organ or gastrointestinal system. Each of the immediately foregoing methods is preferred.

In yet another aspect, this invention provides a method of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective
15 amount of a somatostatin agonist wherein the fibrosis is induced by a disease state and preferably the fibrosis inhibited is in the kidney, lung, liver, skin, central nervous system, bone or bone marrow, cardiovascular system, an endocrine organ or gastrointestinal system. Each of the
20 immediately foregoing methods is preferred.

In yet another aspect, this invention provides a method of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective
25 amount of a somatostatin agonist wherein the fibrosis is induced by an environmental or industrial factor and preferably the fibrosis inhibited is in the kidney, lung, liver, skin, central nervous system, bone or bone marrow, cardiovascular system, an endocrine organ or gastrointestinal system. Each of the immediately foregoing methods is
30 preferred.

In yet another aspect, this invention provides a method of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective amount of a somatostatin agonist wherein the fibrosis is

5


10

15

30

[illegible]

amount of a somatostatin agonist wherein the somatostatin agonist is



10 or a pharmaceutically acceptable salt thereof, wherein

A^1 is a D- or L- isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, β -Nal, β -Pal, Trp, Phe, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe;

15 A^2 is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^3 is pyridyl-Ala, Trp, Phe, β -Nal, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^6 is Val, Ala, Leu, Ile, Nle, Thr, Abu, or Ser;

20 A^7 is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

25 A^8 is a D- or L-isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe;

wherein X for each occurrence is independently selected from the group consisting of CH_3 , Cl, Br, F, OH, OCH_3 and NO_2 ;

30 each R_1 and R_2 , independently, is H, lower acyl or lower alkyl; and R_3 is OH or NH_2 ; provided that at least one of A^1 and A^8 and one of A^2 and A^7 must be an aromatic amino acid; and further provided that A^1 , A^2 , A^7 and A^8 cannot all be aromatic amino acids.

35 In still another aspect, a method of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective amount of a somatostatin agonist

wherein the somatostatin agonist is

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Thr-Phe-Thr-NH₂;

H-D-Phe-p-NO₂-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

H-D-Nal-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

5 H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂;

H-D-Phe-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂; or

H-D-Phe-Ala-Tyr-D-Trp-Lys-Val-Ala-β-D-Nal-NH₂ or a

pharmaceutically acceptable salt thereof.

10 In still another aspect, a method of inhibiting fibrosis in a patient which comprises administering to the patient a therapeutically effective amount of a somatostatin agonist wherein the somatostatin agonist is

D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-β-Nal-NH₂;

15 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Cys-β-Nal-NH₂;

D-β-Nal-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-NH₂;

D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-NH₂;

D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-OH;

20 D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;

Gly-Pen-Phe-D-Trp-Lys-Thr-Cys-Thr-OH;

Phe-Pen-Tyr-D-Trp-Lys-Thr-Cys-Thr-OH;

Phe-Pen-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;

H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-ol;

25 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

H-D-Trp-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

H-D-Trp-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂;

30 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

Ac-D-Phe-Lys'-Tyr-D-Trp-Lys-Val-Asp-Thr-NH₂, wherein an amide bridge is between Lys' and Asp;

Ac-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;


- Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (Bu)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-L-hArg (Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
5 Ac-D-hArg (CH₂CF₃)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt;
Ac-L-hArg (CH₂-CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
10 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me)-Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me)-Thr-Cys-Thr-NHEt;
Ac-hArg (CH₃, hexyl)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
H-hArg (hexyl)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt;
15 Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Propionyl-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys (iPr)-Thr-Cys-Thr-NH₂;
Ac-D-β-Nal-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Gly-hArg (Et)₂-NH₂;
Ac-D-Lys (iPr)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
20 Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Ac-D-hArg (Et)₂-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
25 NH₂;
Ac-Cys-Lys-Asn-4-Cl-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Ser-D-Cys-NH₂;
Bmp-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
Bmp-Tyr-D-Trp-Lys-Val-Cys-Phe-NH₂;
30 Bmp-Tyr-D-Trp-Lys-Val-Cys-p-Cl-Phe-NH₂;
Bmp-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;
H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;

- H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys- β -Nal-NH₂;
H-pentafluoro-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
Ac-D- β -Nal-Cys-pentafluoro-Phe-D-Trp-Lys-Val-Cys-Thr-NH₂;
H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Val-Cys- β -Nal-NH₂;
5 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys- β -Nal-NH₂;
H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
H-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
Ac-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
H-D-Phe-Cys- β -Nal-D-Trp-Lys-Val-Cys-Thr-NH₂;
10 H-D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr-NH₂;
cyclo (Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Thr-N-Me-Phe);
cyclo (N-Me-Ala-Tyr-D-Trp-Lys-Thr-Phe);
15 cyclo (Pro-Tyr-D-Trp-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Thr-Phe);
cyclo (Pro-Phe-L-Trp-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp(F)-Lys-Thr-Phe);
cyclo (Pro-Phe-Trp(F)-Lys-Thr-Phe);
20 cyclo (Pro-Phe-D-Trp-Lys-Ser-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Thr-p-Cl-Phe);
cyclo (D-Ala-N-Me-D-Phe-D-Thr-D-Lys-Trp-D-Phe);
cyclo (D-Ala-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Phe);
cyclo (D-Ala-N-Me-D-Phe-D-Thr-Lys-D-Trp-D-Phe);
25 cyclo (D-Abu-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Tyr);
cyclo (Pro-Tyr-D-Trp-t-4-AchxAla-Thr-Phe);
cyclo (Pro-Phe-D-Trp-t-4-AchxAla-Thr-Phe);
cyclo (N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe);
cyclo (N-Me-Ala-Tyr-D-Trp-t-4-AchxAla-Thr-Phe);
30 cyclo (Pro-Tyr-D-Trp-4-Amphe-Thr-Phe);
cyclo (Pro-Phe-D-Trp-4-Amphe-Thr-Phe);
cyclo (N-Me-Ala-Tyr-D-Trp-4-Amphe-Thr-Phe);
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);

-10-

- cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba-Gaba);
 cyclo (Asn-Phe-D-Trp-Lys-Thr-Phe);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-NH(CH₂)₄CO);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-β-Ala);
 5 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-D-Glu)-OH;
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 10 cyclo (Asn-Phe-Phe-D-Trp(F)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp(NO₂)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-Trp(Br)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe(I)-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Tyr(But)-Gaba);
 15 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-Cys)-OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-Cys)-OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Tpo-Cys)-OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-MeLeu-Cys)-OH;
 20 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Phe-Gaba);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-D-Phe-Gaba);
 cyclo (Phe-Phe-D-Trp(5F)-Lys-Thr-Phe-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys(Ac)-Thr-Phe-NH-(CH₂)₃-CO);
 cyclo (Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 25 cyclo (Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba); or
 cyclo (Orn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba) or a
 pharmaceutically acceptable salt thereof.

In still another aspect, a method of inhibiting fibrosis in a patient which comprises administering to the patient a
 30 therapeutically effective amount of a somatostatin agonist wherein the somatostatin agonist is D-β-Nal-Cys-Tyr-D-Trp-Lys-Thr-Cys-Thr-NH₂, H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂,




or D-Phe-cyclo(Cys-Phe-D-Trp-Lys-Thr-Cys)-Thr-ol or a pharmaceutically acceptable salt thereof. Each of the immediately foregoing methods is preferred.

In a further aspect, this invention provides a method of inhibiting overexpression of TGF- β which comprises administering to a subject an effective amount of somatostatin, somatostatin agonist or a pharmaceutically acceptable salt thereof; preferred of this method is where a somatostatin agonist is administered; a preferred method of the immediately foregoing method is wherein the somatostatin agonist has a higher binding affinity for human somatostatin sub-type receptor 1 than the other human somatostatin sub-type receptors, human somatostatin sub-type receptor 2 than the other human somatostatin sub-type receptors, human somatostatin sub-type receptor 3 than the other human somatostatin sub-type receptors, human somatostatin sub-type receptor 4 than the other human somatostatin sub-type receptors or human somatostatin sub-type receptor 5 than the other human somatostatin sub-type receptors; or wherein the somatostatin agonist has a higher binding affinity for two or more of human somatostatin receptor sub-types e.g., 1, 2, 3, 4 and/or 5. Each of the foregoing methods is preferred.

25 In another further aspect, this invention provides a method of inhibiting overexpression of TGF- β which comprises administering to a subject an effective amount of a somatostatin agonist wherein the somatostatin agonist is

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	

-12-



or a pharmaceutically acceptable salt thereof, wherein

A^1 is a D- or L- isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, β -Nal, β -Pal, Trp, Phe, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe;

A^2 is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^3 is pyridyl-Ala, Trp, Phe, β -Nal, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^6 is Val, Ala, Leu, Ile, Nle, Thr, Abu, or Ser;

A^7 is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^8 is a D- or L-isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe;

wherein X for each occurrence is independently selected from the group consisting of CH₃, Cl, Br, F, OH, OCH₃ and NO₂;

each R_1 and R_2 , independently, is H, lower acyl or lower alkyl; and R_3 is OH or NH₂; provided that at least one of A^1 and A^8 and one of A^2 and A^7 must be an aromatic amino acid; and further provided that A^1 , A^2 , A^7 and A^8 cannot all be aromatic amino acids.

Also, this invention provides a method of inhibiting overexpression of TGF- β which comprises administering to a subject an effective amount of somatostatin agonist wherein the somatostatin agonist is

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Thr-Phe-Thr-NH₂;

H-D-Phe-p-NO₂-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

H-D-Nal-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂;

-13-

H-D-Phe-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;
 H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂; or
 H-D-Phe-Ala-Tyr-D-Trp-Lys-Val-Ala-β-D-Nal-NH₂ or a
 pharmaceutically acceptable salt thereof.

5 Also, this invention provides a method of inhibiting overexpression of TGF-β which comprises administering to a subject an effective amount of somatostatin agonist wherein the somatostatin agonist is

- D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-β-Nal-NH₂;
 10 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Cys-β-Nal-NH₂;
 D-β-Nal-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-NH₂;
 D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-NH₂;
 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-OH;
 15 D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;
 Gly-Pen-Phe-D-Trp-Lys-Thr-Cys-Thr-OH;
 Phe-Pen-Tyr-D-Trp-Lys-Thr-Cys-Thr-OH;
 Phe-Pen-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;
 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-ol;
 20 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 H-D-Trp-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 H-D-Trp-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂;
 25 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 Ac-D-Phe-Lys^{*}-Tyr-D-Trp-Lys-Val-Asp-Thr-NH₂, wherein an amide bridge is between Lys^{*} and Asp;
 Ac-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 30 Ac-D-hArg(Bu)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-L-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(CH₂CF₃)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;


- Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt;
Ac-L-hArg (CH₂-CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
5 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me) -Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me) -Thr-Cys-Thr-NHEt;
Ac-hArg (CH₃, hexyl)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
H-hArg (hexyl₂)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt;
10 Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Propionyl-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys (iPr) -Thr-Cys-Thr-NH₂;
Ac-D-β-Nal-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Gly-hArg (Et)₂-NH₂;
Ac-D-Lys (iPr) -Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
15 Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Ac-D-hArg (Et)₂-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
20 Ac-Cys-Lys-Asn-4-Cl-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Ser-D-Cys-NH₂;
Bmp-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
Bmp-Tyr-D-Trp-Lys-Val-Cys-Phe-NH₂;
25 Bmp-Tyr-D-Trp-Lys-Val-Cys-p-Cl-Phe-NH₂;
Bmp-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;
H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-β-Nal-NH₂;
30 H-pentafluoro-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
Ac-D-β-Nal-Cys-pentafluoro-Phe-D-Trp-Lys-Val-Cys-Thr-NH₂;
H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;


- H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys- β -Nal-NH₂;
H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
H-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
Ac-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
5 H-D-Phe-Cys- β -Nal-D-Trp-Lys-Val-Cys-Thr-NH₂;
H-D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr-NH₂;
cyclo (Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Thr-N-Me-Phe);
10 cyclo (N-Me-Ala-Tyr-D-Trp-Lys-Thr-Phe);
cyclo (Pro-Tyr-D-Trp-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Thr-Phe);
cyclo (Pro-Phe-L-Trp-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp(F)-Lys-Thr-Phe);
15 cyclo (Pro-Phe-Trp(F)-Lys-Thr-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Ser-Phe);
cyclo (Pro-Phe-D-Trp-Lys-Thr-p-Cl-Phe);
cyclo (D-Ala-N-Me-D-Phe-D-Thr-D-Lys-Trp-D-Phe);
cyclo (D-Ala-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Phe);
20 cyclo (D-Ala-N-Me-D-Phe-D-Thr-Lys-D-Trp-D-Phe);
cyclo (D-Abu-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Tyr);
cyclo (Pro-Tyr-D-Trp-t-4-AchxAla-Thr-Phe);
cyclo (Pro-Phe-D-Trp-t-4-AchxAla-Thr-Phe);
cyclo (N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe);
25 cyclo (N-Me-Ala-Tyr-D-Trp-t-4-AchxAla-Thr-Phe);
cyclo (Pro-Tyr-D-Trp-4-Amphe-Thr-Phe);
cyclo (Pro-Phe-D-Trp-4-Amphe-Thr-Phe);
cyclo (N-Me-Ala-Tyr-D-Trp-4-Amphe-Thr-Phe);
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
30 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba-Gaba);
cyclo (Asn-Phe-D-Trp-Lys-Thr-Phe);
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-NH(CH₂)₄CO);
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe- β -Ala);

-16-

- cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-D-Glu) -OH;
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 5 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 cyclo (Asn-Phe-Phe-D-Trp(F)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp(NO₂)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-Trp(Br)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe(I)-Gaba);
 10 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Tyr(But)-Gaba);
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-Cys) -OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-Cys) -OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Tpo-Cys) -OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-MeLeu-Cys) -
 15 OH;
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Phe-Gaba);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-D-Phe-Gaba);
 cyclo (Phe-Phe-D-Trp(5F)-Lys-Thr-Phe-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys(Ac)-Thr-Phe-NH-(CH₂)₃-CO);
 20 cyclo (Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo (Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba); or
 cyclo (Orn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba) or a
 pharmaceutically acceptable salt thereof.

- Also, this invention provides a method of inhibiting
 25 overexpression of TGF- β which comprises administering to a
 subject an effective amount of somatostatin agonist or a
 pharmaceutically acceptable salt thereof wherein the
 somatostatin agonist or a pharmaceutically acceptable salt
 thereof is D- β -Nal-Cys-Tyr-D-Trp-Lys-Thr-Cys-Thr-NH₂, H-Cys-
 30 Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂,





or D-Phe-cyclo(Cys-Phe-D-Trp-Lys-Thr-Cys)-Thr-ol or a pharmaceutically acceptable salt thereof. Each of the foregoing methods is preferred.

5 In still another aspect, this invention provides a method wherein it is preferred that of each of the methods described above that the somatostatin agonist is administered parenterally and more preferably that the somatostatin agonist administered parenterally is administered in a
10 sustained release formulation. It is also preferred that of each of the methods described above that the somatostatin agonist or pharmaceutically acceptable salt thereof is administered orally or topically. Each of the foregoing methods is preferred.

15 Still another aspect of the present invention provides a pharmaceutical composition useful for inhibiting fibrosis in a patient which comprises a pharmaceutically acceptable carrier and an effective amount of somatostatin, somatostatin agonist or a pharmaceutically acceptable salt thereof,
20 preferred of the immediately foregoing pharmaceutical composition is a pharmaceutical composition which comprises a somatostatin agonist or a pharmaceutically acceptable salt thereof.

Still another aspect of the present invention provides a
25 pharmaceutical composition useful for inhibiting overexpression of TGF- β which comprises a pharmaceutically acceptable carrier and an effective amount of somatostatin, somatostatin agonist or a pharmaceutically acceptable salt thereof, preferred of the immediately foregoing
30 pharmaceutical composition is a pharmaceutical composition which comprises a somatostatin agonist or a pharmaceutically

acceptable salt thereof.

Definition of "somatostatin agonist" will be defined below. A therapeutically effective amount depends upon the condition being treated, treatment regimen, the route of administration chosen, and the specific activity of the compound used and ultimately will be decided by the attending physician or veterinarian. In one embodiment, the somatostatin agonist is administered to the patient until the fibrotic process is arrested and/or is reversed. In another embodiment, the somatostatin agonist is administered for the lifetime of the patient. In still another embodiment, the somatostatin agonist is administered prior to the event which initiates the fibrotic process (e.g., prior to chemotherapy or exposure to radiation such as in radiation therapy).

Somatostatin or a somatostatin agonist may be injected parenterally, e.g., intravenously, into the bloodstream of the subject being treated. However, it will be readily appreciated by those skilled in the art that the route, such as subcutaneous, intramuscular, intraperitoneal, enterally, transdermally, transmucously, sustained released polymer compositions (e.g., a lactic acid polymer or lactic-glycolic acid copolymer microparticle or implant), perfusion, nasal, oral, topical, vaginal, rectal, nasal, sublingual, etc., will vary with the condition being treated and the activity and bioavailability of the somatostatin agonist being used.

The dosage of active ingredient administered in a method of this invention may be varied; however, it is necessary that the amount of the active ingredient be such that a suitable dosage form is obtained. The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment. Generally, dosage levels of between 0.000001 to 100 mg/kg of body weight daily are administered to humans and other animals, e.g., mammals.

A preferred dosage range is 0.01 to 5.0 mg/kg of body weight daily which can be administered as a single dose or divided into multiple doses.

While it is possible for the somatostatin agonist to be administered as the pure or substantially pure compound, it may also be presented as a pharmaceutical formulation or preparation. The formulations to be used in the present invention, for both humans and animals, comprise any of the somatostatin agonists to be described below, together with one or more pharmaceutically acceptable carriers thereof, and optionally other therapeutic ingredients.

The carrier must be "acceptable" in the sense of being compatible with the active ingredient(s) of the formulation (e.g., capable of stabilizing peptides) and not deleterious to the subject to be treated. Desirably, the formulation should not include oxidizing agents or other substances with which peptides are known to be incompatible. For example, somatostatin agonists in the cyclized form (e.g., internal cysteine disulfide bond) are oxidized; thus, the presence of reducing agents as excipients could lead to an opening of the cysteine disulfide bridge. On the other hand, highly oxidative conditions can lead to the formation of cysteine sulfoxide and to the oxidation of tryptophane. Consequently, it is important to carefully select the excipient. pH is another key factor, and it may be necessary to buffer the product under slightly acidic conditions (pH 5 to 6).

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient(s) into association with the carrier which constitutes one or more accessory ingredients.

In general, the formulations for tablets or powders are prepared by uniformly and intimately blending the active

-20-

ingredient with finely divided solid carriers, and then, if necessary, as in the case of tablets, forming the product into the desired shape and size.

Formulations suitable for parenteral (e.g., intravenous) administration, on the other hand, conveniently comprise sterile aqueous solutions of the active ingredient(s). Preferably, the solutions are isotonic with the blood of the subject to be treated. Such formulations may be conveniently prepared by dissolving active ingredient(s) in a solvent comprising water to produce an aqueous solution, and rendering said solution sterile. The formulation may be presented in unit or multi-dose containers, for example, sealed ampoules or vials.

Formulations suitable for sustained release parenteral administrations (e.g., biodegradable polymer formulations) are also well known in the art. See, e.g., U.S. Patent Nos. 3,773,919 and 4,767,628, the teachings of which are incorporated herein by reference, and PCT Publication No. WO 94/15587.

Compositions for rectal or vaginal administration are preferably suppositories which may contain, in addition to the active substance, excipients such as cocoa butter or a suppository wax.

Compositions for nasal or sublingual administration are also prepared with standard excipients well known in the art.

For topical administration, they are best used in the form of solutions, creams, salves, lotions, ointments and the like.

The somatostatin or somatostatin agonist may also be administered with known initiators (e.g., chemotherapeutics) of the fibrotic process to ameliorate fibrosis or to prevent the initiation of fibrosis.

Other features and advantages of the invention will be apparent from the following description of the preferred

embodiments and from the claims.

Abbreviations

β -Nal = β -naphthylalanine

β -Pal = β -pyridylalanine

5 hArg(Bu) = N-guanidino-(butyl)-homoarginine

hArg(Et)₂ = N,N'-guanidino-(diethyl)-homoarginine

hArg(CH₂CF₃)₂ = N,N'-guanidino-bis-(2,2,2,-trifluoroethyl)-
homoarginine

hArg(CH₃, hexyl) = N,N'-guanidino-(methyl, hexyl)-homoarginine

10 Lys(Me) = N^ε-methyllysine

Lys(iPr) = N^ε-isopropyllysine

AmPhe = aminomethylphenylalanine

AChxAla = aminocyclohexylalanine

Abu = α -aminobutyric acid

15 Tpo = 4-thiaproline

MeLeu = N-methylleucine

Orn = ornithine

Nle = norleucine

Nva = norvaline

20 Trp(Br) = 5-bromo-tryptophan

Trp(F) = 5-fluoro-tryptophan

Trp(NO₂) = 5-nitro-tryptophan

Gaba = γ -aminobutyric acid

Bmp = β -mercaptopropionyl

25 Ac = acetyl

Pen = pencillamine.

DETAILED DESCRIPTION OF THE INVENTION

30 It is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent. The following specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way

whatsoever.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Also, all publications, patent applications, patents, and other references mentioned herein are incorporated by reference.

The fibrosis which is inhibited can be located in various parts of the body and can be of a particular kind, for example, the fibrosis may be located:

in the kidney, for example, fibrosis as observed in glomerulonephritis (see Yoshioka et al., Lab Invest 1993;68: 154-63), diabetic nephropathy (see Yamamoto et al., Proc Natl Acad Sci USA 1993;90: 1814-8), allograft rejection (see Shihab et al., J Am Soc Nephrol 1993;4: 671, abstract), and HIV nephropathy (see Border et al., J Am Soc Nephrol 1993;4: 675, abstract);

in the liver, for example, cirrhosis (see Castilla et al., N Engl J Med 1991;324: 933-940 and Nagy et al., Hepatology 1991;14: 269-73), and veno-occlusive disease (see Anscher et al., N Engl J Med 1993;328: 1592-8);

in the lung, for example, idiopathic fibrosis (see Anscher et al., N Engl J Med 1993;328: 1592-8 and Brockelmann et al., Proc Natl Acad Sci USA 1991;88: 6642-6) and autoimmune fibrosis (see Deguchi, Ann Rheum Dis 1992;51: 362-5);

in the skin, for example, systemic sclerosis (see Kulozik et al., J Clin Invest 1990;86: 917-22), keloids (see Peltonen et al., J Invest Dermatol 1991;97: 240-8), scars (see Ghahary et al., J Lab Clin Med 1993;122: 465-73), and eosinophilia-myalgia syndrome (see Varga et al., Ann Intern Med 1992;116: 140-7);

in the central nervous system, for example, intraocular fibrosis (see Conner et al., J Clin Invest 1989;83: 1661-6);

-23-

in the cardiovascular system, for example, vascular restenosis (see Nikol et al., J Clin Invest 1992;90: 1582-92);

in the nose, for example, nasal polyposis (see Ohno et al., J Clin Invest 1992;89: 1662-8);

in bone or bone marrow (see Harrison's Principles of Internal Medicine, Thirteenth Edition, Volume 2, Chapter 362, pp. 2197-2199; Najean, Y. et al., Leuk Lymphoma, 1996, 22 Suppl 1:111-119; and Reith, J.D. et al., Am J Surg Pathol, 1996 20(11): 1368-1377);

in an endocrine organ (see Endocrinology, Third Edition, Edited by Leslie J. DeGroot, Vol. 1, pp. 165-177 and pp. 747-751);

and in the gastrointestinal system (see Mizoi, T. et al, Cancer Res., 1993 53(1): 183-190; and Tahara, E., J. Cancer Res. Clin. Oncol., 1990, 116(2), 121-131).

A fibrotic disorder may be induced by a number of causes including:

chemotherapy, for example, pulmonary fibrosis resulting from bleomycin, chlorambucil, cyclophosphamide, methotrexate, mustine, or procarbazine treatment (see Key Facts in Oncology by Lilly, Drug Therapy, p.11, 1994);

radiation exposure whether accidental or purposeful as in radiation therapy, for example, interstitial lung disease (ILD) resulting from radiation (see Cecil Textbook of Medicine, 19th Edition, edited by James B. Wyngaarden, Lloyd H. Smith, Jr., and J. Claude Bennet, Chapter 60, Table 60-5, p. 399, 1992);

environmental or industrial factors or pollutants such as chemicals, fumes, metals, vapors, gases, etc., for example, ILD resulting from asbestos or coal dust (see Cecil Textbook of Medicine, 19th Edition, edited by James B. Wyngaarden, Lloyd H. Smith, Jr., and J. Claude Bennet, Chapter 60, Table 60-2, p. 398, 1992);

-24-

a drug or a combination of drugs, for example, antibiotics (e.g. penicillins, sulfonamides, etc.), cardiovascular drugs (e.g. hydralazine, beta blockers, etc.), CNS drugs (phenytoin, chlorpromazine, etc.) anti-inflammatory
5 drugs (e.g. gold salts, phenylbutazone, etc.), etc. can cause
ILD (see Cecil Textbook of Medicine, 19th Edition, edited by James B. Wyngaarden, Lloyd H. Smith, Jr., and J. Claude Bennet, Chapter 60, Table 60-4, p. 398, 1992);

an immune reaction disorder, for example, chronic graft-
10 vs-host disease with dermal fibrosis, (see Fibrotic Skin Diseases, Editorial, J. Uitto and S. Jimenez, Arch, Dermatol, Vol 126, May 1990, p.662);

disease states such as aspiration pneumonia which is a known cause of ILD, (see Harrison's Principles of Internal
15 Medicine, Twelfth Edition, Chapter 211, Table 211-1, P 1083) and parasite induced fibrosis (see Wahl, S.M., Kidney Int, 1997, 51(5): 1370-1375); and

wounds, for example, blunt trauma, surgical incisions, battlefield wounds, etc., as in penetrating injuries of the
20 CNS (see Ann Logan, et al., Brain Research, 587 (1992), 216-225).

Somatostatin and its Agonists

Somatostatin (somatotropin release inhibiting factor or SRIF) has both a 14 amino acid isoform (somatostatin-14) and
25 a 28 amino acid isoform (somatostatin-28). See Wilson, J. & Foster, D., *Williams Textbook of Endocrinology*, p. 510 (7th ed., 1985). The compound is an inhibitor of secretion of the growth hormone and was originally isolated from the hypothalamus. Brazeau et al., *Science* 179:77 (1973). Native
30 somatostatin has a very short duration of effect in vivo since it is rapidly inactivated by endo- and exopeptidase. Many novel analogs have been prepared in order to enhance the duration of effect, biological activity, and selectivity (e.g., for the particular somatostatin receptor) of this

-25-

hormone. Such analogs will be called "somatostatin agonists" herein. Further, compounds that are short peptides modified by organic moieties and non-peptides, such as organic molecules that do not have an art-recognized amino acid as part of its structure, that bind to somatostatin receptor(s) are also within the meaning of "somatostatin agonists".

Various somatostatin receptors (SSTRs) have been isolated, e.g., SSTR-1, SSTR-2, SSTR-3, SSTR-4, and SSTR-5. Thus, the somatostatin agonist may be a SSTR-1 agonist, SSTR-2 agonist, SSTR-3 agonist, SSTR-4 agonist or a SSTR-5 agonist. In one embodiment, the somatostatin agonist is an SSTR-2 agonist or an SSTR-5 agonist. What is meant by an "SSTR-2 agonist" or an "SSTR-5 agonist" is a compound which (1) has a high affinity (e.g., K_i of less than 1 nM or, preferably, of less than 10 nM) for the SSTR-2 or SSTR-5, respectively (as defined by the receptor binding assay described below), and (2) inhibits the formation of fibrosis (e.g., as defined by the biological assay described below). The somatostatin agonist may also be selective for a particular somatostatin receptor, e.g., have a higher binding affinity for a particular somatostatin receptor subtype. In one embodiment, the somatostatin receptor is an SSTR-2 or SSTR-5 selective agonist.

Somatostatin agonists which can be used to practice the therapeutic method of the present invention include, but are not limited to, those covered by formulae or those specifically recited in the publications set forth below, all of which are hereby incorporated by reference.

EP Application No. P5 164 EU (Inventor: G. Keri);
Van Binst, G. et al. Peptide Research 5:8 (1992);
Horvath, A. et al. Abstract, "Conformations of Somatostatin Analogs Having Antitumor Activity", 22nd European peptide Symposium, September 13-19, 1992,

Interlaken, Switzerland:

- PCT Application WO 91/09056 (1991);
EP Application 0 363 589 A2 (1990);
U.S. Patent No. 4,904,642 (1990);
5 U.S. Patent No. 4,871,717 (1989);
U.S. Patent No. 4,853,371 (1989);
U.S. Patent No. 4,725,577 (1988);
U.S. Patent No. 4,684,620 (1987);
U.S. Patent No. 4,650,787 (1987);
10 U.S. Patent No. 4,603,120 (1986);
U.S. Patent No. 4,585,755 (1986);
EP Application 0 203 031 A2 (1986);
U.S. Patent No. 4,522,813 (1985);
U.S. Patent No. 4,486,415 (1984);
15 U.S. Patent No. 4,485,101 (1984);
U.S. Patent No. 4,435,385 (1984);
U.S. Patent No. 4,395,403 (1983);
U.S. Patent No. 4,369,179 (1983);
U.S. Patent No. 4,360,516 (1982);
20 U.S. Patent No. 4,358,439 (1982);
U.S. Patent No. 4,328,214 (1982);
U.S. Patent No. 4,316,890 (1982);
U.S. Patent No. 4,310,518 (1982);
U.S. Patent No. 4,291,022 (1981);
25 U.S. Patent No. 4,238,481 (1980);
U.S. Patent No. 4,235,886 (1980);
U.S. Patent No. 4,224,190 (1980);
U.S. Patent No. 4,211,693 (1980);
U.S. Patent No. 4,190,648 (1980);
30 U.S. Patent No. 4,146,612 (1979); and
U.S. Patent No. 4,133,782 (1979).

Examples of somatostatin agonists include, but are not limited to, the following somatostatin analogs and pharmaceutically acceptable salt thereof which are disclosed

in the above-cited references:

- D- β -Nal-Cys-Tyr-D-Trp-Lys-Thr-Cys-Thr-NH₂ (BIM-23014);
 D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys- β -Nal-NH₂;
 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Cys- β -Nal-NH₂;
 5 D- β -Nal-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-NH₂;
 D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-NH₂;
 D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-OH;
 D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;
 10 Gly-Pen-Phe-D-Trp-Lys-Thr-Cys-Thr-OH;
 Phe-Pen-Tyr-D-Trp-Lys-Thr-Cys-Thr-OH;
 Phe-Pen-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;
 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-ol;
 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 15 H-D-Trp-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 H-D-Trp-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 20 Ac-D-Phe-Lys-Tyr-D-Trp-Lys-Val-Asp-Thr-NH₂ (an amide
 bridge formed between Lys and Asp);
 Ac-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(Bu)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 25 Ac-D-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-L-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(CH₂CF₃)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
 Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-
 NH₂;
 30 Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-
 NH₂;
 Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-
 NH₂;

-28-

Ac-L-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys(Me)-Thr-Cys-Thr-NH₂;

5 Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys(Me)-Thr-Cys-Thr-NHEt;

Ac-hArg(CH₃, hexyl)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

10 H-hArg(hexyl)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt;
Ac-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;
Propionyl-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys(iPr)-Thr-Cys-Thr-NH₂;

15 Ac-D-β-Nal-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Gly-hArg(Et)₂-NH₂;

Ac-D-Lys(iPr)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;
Ac-D-hArg(CH₂CF₃)₂-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

20 Ac-D-hArg(CH₂CF₃)₂-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;

Ac-D-hArg(Et)₂-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-Cys-Lys-Asn-4-Cl-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Ser-D-Cys-NH₂;

25 Bmp-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

Bmp-Tyr-D-Trp-Lys-Val-Cys-Phe-NH₂;

Bmp-Tyr-D-Trp-Lys-Val-Cys-p-Cl-Phe-NH₂;

Bmp-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;

H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

30 H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-β-Nal-NH₂;

H-pentafluoro-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

Ac-D-β-Nal-Cys-pentafluoro-Phe-D-Trp-Lys-Val-Cys-Thr-

NH₂;

- H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;
- H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;
- H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
- 5 H-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
- Ac-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
- H-D-Phe-Cys-β-Nal-D-Trp-Lys-Val-Cys-Thr-NH₂;
- H-D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr-NH₂;
- cyclo (Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe) ;
- 10 cyclo (Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe) ;
- cyclo (Pro-Phe-D-Trp-Lys-Thr-N-Me-Phe) ;
- cyclo (N-Me-Ala-Tyr-D-Trp-Lys-Thr-Phe) ;
- cyclo (Pro-Tyr-D-Trp-Lys-Thr-Phe) ;
- cyclo (Pro-Phe-D-Trp-Lys-Thr-Phe) ;
- 15 cyclo (Pro-Phe-L-Trp-Lys-Thr-Phe) ;
- cyclo (Pro-Phe-D-Trp(F)-Lys-Thr-Phe) ;
- cyclo (Pro-Phe-Trp(F)-Lys-Thr-Phe) ;
- cyclo (Pro-Phe-D-Trp-Lys-Ser-Phe) ;
- cyclo (Pro-Phe-D-Trp-Lys-Thr-p-Cl-Phe) ;
- 20 cyclo (D-Ala-N-Me-D-Phe-D-Thr-D-Lys-Trp-D-Phe) ;
- cyclo (D-Ala-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Phe) ;
- cyclo (D-Ala-N-Me-D-Phe-D-Thr-Lys-D-Trp-D-Phe) ;
- cyclo (D-Abu-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Tyr) ;
- cyclo (Pro-Tyr-D-Trp-t-4-AchxAla-Thr-Phe) ;
- 25 cyclo (Pro-Phe-D-Trp-t-4-AchxAla-Thr-Phe) ;
- cyclo (N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe) ;
- cyclo (N-Me-Ala-Tyr-D-Trp-t-4-AchxAla-Thr-Phe) ;
- cyclo (Pro-Tyr-D-Trp-4-Amphe-Thr-Phe) ;
- cyclo (Pro-Phe-D-Trp-4-Amphe-Thr-Phe) ;
- 30 cyclo (N-Me-Ala-Tyr-D-Trp-4-Amphe-Thr-Phe) ;
- cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba) ;
- cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba-Gaba) ;
- cyclo (Asn-Phe-D-Trp-Lys-Thr-Phe) ;

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058

-30-


- cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-NH(CH₂)₃CO);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-β-Ala);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-D-Glu)-OH;
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe);
 5 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 cyclo (Asn-Phe-Phe-D-Trp(F)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp(NO₂)-Lys-Thr-Phe-Gaba);
 10 cyclo (Asn-Phe-Phe-Trp(Br)-Lys-Thr-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe(I)-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Tyr(But)-Gaba);
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-
 Cys)-OH;
 15 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-
 Cys)-OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Tpo-
 Cys)-OH;
 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-
 20 MeLeu-Cys)-OH;
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Phe-Gaba);
 cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-D-Phe-Gaba);
 cyclo (Phe-Phe-D-Trp(5F)-Lys-Thr-Phe-Phe-Gaba);
 cyclo (Asn-Phe-Phe-D-Trp-Lys(Ac)-Thr-Phe-NH-(CH₂)₃-
 25 CO);
 cyclo (Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo (Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo (Orn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba); and
 H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂(BIM-23268).
 30 Note that for all somatostatin agonists described
 herein, each amino acid residue represents the structure of
 -NH-C(R)H-CO-, in which R is the side chain (e.g., CH₃ for
 Ala) except for Thr-ol which means -NH-CH(CH(CH₃)OH)-CH₂-OH

-3i-

and Pro which means prolinyl. Lines between amino acid residues represent peptide bonds which join the amino acids.

Also, where the amino acid residue is optically active, it is the L-form configuration that is intended unless D-form is expressly designated. A disulfide bridge is formed between two Cys residues; however, it is not shown.

Use of linear somatostatin agonists of the following formula is also within the invention:



or a pharmaceutically acceptable salt thereof, wherein

A^1 is a D- or L- isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, β -Nal, β -Pal, Trp, Phe, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe;

A^2 is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^3 is pyridyl-Ala, Trp, Phe, β -Nal, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^6 is Val, Ala, Leu, Ile, Nle, Thr, Abu, or Ser;

A^7 is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe;

A^8 is a D- or L-isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe;

each R_1 and R_2 , independently, is H, lower acyl or lower alkyl; and R_3 is OH or NH_2 ; provided that at least one of A^1 and A^8 and one of A^2 and A^7 must be an aromatic amino acid; and further provided that A^1 , A^2 , A^7 and A^8 cannot all be aromatic amino acids.

Examples of linear agonists to be used in the method of this invention include:

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Thr-Phe-Thr-NH₂;

H-D-Phe-p-NO₂-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

5 H-D-Nal-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂;


H-D-Phe-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;


and

10 H-D-Phe-Ala-Tyr-D-Trp-Lys-Val-Ala-β-D-Nal-NH₂ or a pharmaceutically acceptable salt thereof.



If desired, one or more chemical moieties, e.g., a sugar derivative, mono- or poly-hydroxy C₂₋₁₂ alkyl, mono or poly-hydroxy C₂₋₁₂ acyl groups, or a piperazine derivative, can be attached to the somatostatin agonist, e.g., to the N-terminus amino acid. See PCT Application WO 88/02756, European Application 0 329 295, and PCT Application No. WO 94/04752. An example of a somatostatin agonists which contain N-terminal chemical substitutions are:



;



;



(BIM-23197) or a pharmaceutically acceptable salt thereof.

Synthesis of somatostatin agonists

The methods for synthesizing somatostatin agonists are well documented and are within the ability of a person of ordinary skill in the art, for example, as illustrated in the U.S. Patents and other references cited hereinabove.

Synthesis of short amino acid sequences is well established in the peptide art. For example, synthesis of D-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂, described above, can be synthesized by following the protocol set forth in U.S. Patent No. 4,853,371 and synthesis of H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂, described above, can be achieved by following the protocol set forth in Example I of European Patent Application 0 395 417 A1. The synthesis of somatostatin agonists with a substituted N-terminus can be achieved, for example, by following the protocol set forth in WO 88/02756, European Patent Application No. 0 329 295, and PCT Publication No. WO 94/04752.

Somatostatin Receptor Binding Assays

The human SSTR-1, SSTR-2, SSTR-3, SSTR-4, and SSTR-5 cDNA clones have been described (SSTR-1 and SSTR-2 in Yamada, Y., et al., Proc. Natl. Acad. Sci. USA., 89:251-255 (1992); SSTR-3 in Yamada, et al., Mol. Endocrinol. 6:2136-2142 (1993); and SSTR-4 and SSTR-5 in Yamada, et al., Biochem. Biophys. Res. Commun. 195:844-852 (1993)) and are also available from American Type Culture Collection (ATCC, Rockville, MD) (ATCC Nos. 79044 (SSTR-1), 79046 (SSTR-2), and 79048 (SSTR-3)). Based on the restriction endonuclease maps, the entire coding region of each SSTR cDNA may be excised by suitable restriction endonuclease digestion (Maniatis, T., et al., *Molecular Cloning - A Laboratory Manual*, CSHL, 1982). Restriction endonucleases are available from New England Biolabs (Beverly, MA). This cDNA fragment was inserted into the mammalian expression vector, pCMV (Russell, D., et al.,

-34-

J. Biol. Chem., 264:8222-8229 (1989)), using standard molecular biology techniques (see e.g., Maniatis, T., et al., Molecular Cloning, -A Laboratory Manual, Cold Spring Harbor Laboratory, 1982) to produce the expression plasmid, pCMV-human SSTR-1 through pCMV-human SSTR-5. Other mammalian expression vectors include pcDNA1/Amp (Invitrogen, Sandlesy, CA). The expression plasmids were introduced into the suitable bacterial host, E. Coli HB101 (Stratagene, La Jolla, CA) and plasmid DNAs, for transfection, were prepared on Cesium Chloride gradients.

CHO-K1 (ovary, Chinese hamster) cells were obtained from ATCC (ATCC No. CCL 61). The cells were grown and maintained in Ham's F12 media (Gibco BRL, Grand Island, NY) supplemented with 10% fetal bovine serum under standard tissue culture conditions. For transfection, the cells were seeded at a density 1×10^6 /60-cm plate (Baxter Scientific Products, McGaw Park, IL.). DNA mediated transfection was carried out using the calcium phosphate co-precipitation method (Ausubel, F.M., et al., Current Protocols in Molecular Biology, John Wiley & Sons, 1987). The plasmid pRSV-neo (ATCC; ATCC No. 37198) was included as a selectable marker at 1/10 the concentration of the expression plasmid. CHO-K1 clonal cell lines that have stably inherited the transfected DNA were selected for growth in Ham's F12 media containing 10% fetal bovine serum and 0.5mg/ml of G418 (Sigma). The cells were ring-cloned and expanded in the same media for analysis.

Expression of the human SSTR-1 through SSTR-5 receptors in the CHO-K1 cells were detected by Northern blot analysis of total RNA prepared from the cells (Sambrook, J.E., et al., Molecular Cloning - A Laboratory Manual, Ed. 2., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989) and by receptor binding using [125 I-Tyr 11]somatostatin-14 as a ligand.

Transfected cell lines expressing the human SSTR receptors were clonally expanded in culture and used in the following

SSTR binding protocol.

Crude membranes were prepared by homogenization of the transfected cells in 20 ml of ice-cold 50 mM Tris-HCl with a tissue homogenizer (setting 6, 15 sec). Buffer was added to
5 obtain a final volume of 40 ml, and the homogenate was centrifuged in a Sorval® SS-34 rotor (Sorval, Newtown, Connecticut) at 39,000 g for 10 min at 0-4°C. The resulting supernatant was decanted and discarded. The pellet was rehomogenized in ice-cold buffer, diluted, and centrifuged as
10 before. The final pellet was resuspended in the 10 mM Tris HCl and held on ice for the receptor binding assay.

Aliquots of the membrane preparation were incubated for 30 min at 30°C with 0.05 nM [¹²⁵I-Tyr¹¹]somatostatin-14 (2000 Ci/mmol; Amersham Corp., Arlington Heights, IL) in 50 mM
15 HEPES (pH 7.4) containing a test somatostatin agonist of various concentrations (e.g., 10⁻¹¹ to 10⁻⁶), 10 mg/ml bovine serum albumin (fraction V) (Sigma Chemical Co., St. Louis, MO), MgCl₂ (5 mM), Trasylol (also known as aprotinin) (Sigma Chemical Co.) (200 KIU ml), bacitracin (Sigma Chemical Co.)
20 (0.02 mg/ml), and phenylmethylsulphonyl fluoride (Sigma Chemical Co.) (0.02 mg/ml). The final assay volume was 0.3 ml. The incubations were terminated by rapid filtration through GF/C filters (pre-soaked in 0.3% polyethylenimine for 30 min) using a Brandel filtration manifold (Brandel Research
25 and Development Co., Gaithersburg, Maryland). Each tube and filter were then washed three times with 5 ml aliquots of ice-cold buffer. Specific binding was defined as the total [¹²⁵I-Tyr¹¹]somatostatin-14 bound minus that bound in the presence of 1000 nM of somatostatin-14. The K_i values for
30 the tested somatostatin agonists were calculated by using the following formula: $K_i = IC_{50} / [1 + (LC/LEC)]$ where IC₅₀ is the concentration of test somatostatin agonist required to inhibit 50 percent of the specific binding of the radioligand

-36-

[¹²⁵I-Tyr¹¹]somatostatin-14, LC is the concentration of the radioligand (0.05 nM), and LEC is the equilibrium dissociation constant of the radioligand (0.16 nM). The K_i values (nM) for the tested somatostatin agonists are shown in Table I.

TABLE I

	hSSTR-1	hSSTR-2	hSSTR-3	hSSTR-4	hSSTR-5
Somatostatin-14	2.256	0.71	1.432	1.768	0.883
Somatostatin-28	2.382	0.57	1.021	7.93	0.383
BIM-23014	2414	1.10	121	1826	5.21
BIM-23190	5210	0.47	2154	7537	11.1
BIM-23197	6016	0.09	26.8	3897	9.81
BIM-23268	12.27	6.84	62	19.96	0.38

Inhibition of Fibrosis

The somatostatin agonists may be tested for their ability to inhibit fibrosis.

(a) Demonstration of Anti-Fibrotic Activity *In Vitro*

Rats are injected either with anti-thymocyte serum (ATS) (see S. Okuda et al., J. Clin. Invest., Vol. 86, 1990, pp. 453-462) to induce glomerulonephritis or with phosphate buffered saline (PBS) to serve as controls. Six days later, the kidneys are removed, and the glomeruli are isolated and placed in culture for 72 hours. Culture conditions consist of 2000 glomeruli/well in a 1 ml volume of serum-free RPMI 1640 (with insulin supplementation) (Gibco, Gaithersburg, Maryland). Test somatostatin or somatostatin agonists are added at the time of culture. The supernatant from the cultures is collected and stored at -70°C until assayed to determine the concentration of collagen I, transforming

growth factor β -1 (TGF β -1), fibronectin containing an extra domain A (fibronectin EDA+), and plasminogen activator inhibitor I (PAI-I) as markers of fibrotic activity. In addition, individual glomeruli are examined by immunofluorescent staining and scored for relevant matrix proteins. Values were compared between PBS-treated, negative fibrotic control glomeruli; ATS-treated, non-drug treated, positive fibrotic control glomeruli; and the ATS-treated, drug treated, fibrotic glomeruli to determine the degree to which the fibrotic process is inhibited by somatostatin or the somatostatin agonists.

(b) Demonstration of Anti-Fibrotic Activity *In Vivo*

Rats are injected either with anti-thymocyte serum (ATS) to induce glomerulonephritis or with phosphate buffered saline (PBS) as a control. One hour later, treatment is initiated with somatostatin or a somatostatin agonist. Somatostatin or the somatostatin agonist are administered subcutaneously twice per day for 5 days. On day 5 the rats are placed in metabolic cages, and 24 hour urine is collected to determine protein content. On day 6, the kidneys are removed, and tissue samples are either placed in formalin or frozen for histological evaluation. Glomeruli are isolated from the remaining tissue and are placed in culture for 72 hours. Culture conditions consisted of 2000 glomeruli/well in a 1 ml volume of serum free RPMI 1640 (with insulin supplementation). The supernatant from the cultures are collected and stored at -70°C until assayed to determine the concentration of collagen I, transforming growth factor β -1 (TGF β -1), fibronectin containing an extra domain A (fibronectin EDA+), and plasminogen activator inhibitor I (PAI1) as markers of fibrotic activity. The presence of matrix proteins is measured via immunofluorescent staining of frozen kidney sections with antibodies to matrix proteins

-38-

induced by TGF β -1 such as fibronectin EDA+, collagen I, PAI1, and tenascin. From the cultured isolated glomeruli direct measurements of TGF β -1, PAI1, and fibronectin secreted into the culture supernatant can be determined via ELISAs (enzyme-linked immunoabsorbent assay). Glomeruli from samples in each group can be used to extract mRNA and the message levels for TGF β -1, GADPH, collagen I, collagen III, fibronectin, and PAI1 determined by Northern analysis. As an indicator of gross histological changes, PAS (periodic acid-Schiff) stained paraffin sections are graded on the basis of their pathological matrix scores. Values are compared between PBS-treated, negative fibrotic control animals; ATS-treated, non-drug treated, positive fibrotic control animals; and the ATS-treated, drug-treated animals to determine the degree to which the fibrotic process is inhibited by somatostatin or the somatostatin agonist.

OTHER EMBODIMENTS

The foregoing description has been limited to specific embodiments of this invention. It will be apparent, however, that variations and modifications may be made to the invention, with the attainment of some or all of the advantages of the invention. Such embodiments are also within the scope of the following claims.